

Interactive effects of temperature and drought on cassava growth and toxicity: implications for food security?

ALICIA L. BROWN, TIMOTHY R. CAVAGNARO*, ROS GLEADOW and REBECCA E. MILLER†

School of Biological Sciences, Monash University, Clayton, Vic. 3800, Australia

Abstract

Cassava is an important dietary component for over 1 billion people, and its ability to yield under drought has led to it being promoted as an important crop for food security under climate change. Despite its known photosynthetic plasticity in response to temperature, little is known about how temperature affects plant toxicity or about interactions between temperature and drought, which is important because cassava tissues contain high levels of toxic cyanogenic glucosides, a major health and food safety concern. In a controlled glasshouse experiment, plants were grown at 2 daytime temperatures (23 °C and 34 °C), and either well-watered or subject to a 1 month drought prior to harvest at 6 months. The objective was to determine the separate and interactive effects of temperature and drought on growth and toxicity. Both temperature and drought affected cassava physiology and chemistry. While temperature alone drove differences in plant height and above-ground biomass, drought and temperature × drought interactions most affected tuber yield, as well as foliar and tuber chemistry, including C : N, nitrogen and cyanide potential (CN_p; total cyanide released from cyanogenic glucosides). Conditions that most stimulated growth and yield (well-watered × high temperature) effected a reduction in tuber toxicity, whereas drought inhibited growth and yield, and was associated with increased foliar and tuber toxicity. The magnitude of drought effects on tuber yield and toxicity were greater at high temperature; thus, increases in tuber CN_p were not merely a consequence of reduced tuber biomass. Findings confirm that cassava is adaptable to forecast temperature increases, particularly in areas of adequate or increasing rainfall; however, in regions forecast for increased incidence of drought, the effects of drought on both food quality (tuber toxicity) and yield are a greater threat to future food security and indicate an increasing necessity for processing of cassava to reduce toxicity.

Keywords: climate change, cyanogenesis, food security, *Manihot esculenta*, mycorrhizas, nitrogen, nutrition, plant defence

Received 17 December 2015; revised version received 22 April 2016 and accepted 4 May 2016

Introduction

Cassava, *Manihot esculenta* Crantz, is eaten by approximately one billion people every day, mainly in the tropical and subtropical regions of Asia, Latin America and Africa and is the major staple for 35–50% of people living in different areas of sub-Saharan Africa (FAO & IFAD, 2000). Cassava grows in a wide range of soil and climatic conditions, is easily propagated, resistant to drought and pests, and the tuberous roots have extended viability (up to 3 years) when left in soil (Nhassico *et al.*, 2008). These traits help explain the sizeable increase in the proportion of cultivated land taken up by cassava in Africa since the 1970s (Fermont

et al., 2008) and underpin its promotion as increasingly important for food security, particularly in the context of climate change (e.g. Jarvis *et al.*, 2012).

Although cassava yields well under poor conditions, the tuberous roots are low in nutritional quality – an important component of food security (Pinstrup-Andersen, 2009). Tubers are high in carbohydrate (80–90% dry matter), but low in protein (1–3% dry matter), low in micronutrients and contain cyanogenic glucosides (Montagnac *et al.*, 2009). Cyanogenic glucosides, which are produced primarily as a defence against herbivores, are hydrolysed to release toxic hydrogen cyanide when the leaves and tubers are crushed or chewed (Conn, 1981). Consumption of cassava-based food stuffs that are inadequately processed to remove cyanogenic compounds can cause acute poisoning resulting in headaches and vomiting, and may lead to a type of permanent leg paralysis known as Konzo, or even death (Cliff, 1994). Increasing penetration of these products into communities without understanding the risks also poses a potential health hazard (Burns *et al.*, 2012a). Critically, cyanogenic glucoside

*Present address: Waite Research Institute, School of Agriculture, Food and Wine, The University of Adelaide, PMB1, Glen Osmond, SA 5064, Australia.

†Present address: School of Ecosystem and Forest Sciences, University of Melbourne, Burnley Campus, Richmond, Vic. 3121, Australia.

Correspondence: Rebecca E. Miller, tel. +61 3 9035 6995, fax +61 3 9035 6800, e-mail: miller.r@unimelb.edu.au

concentrations in cassava vary with genotype and climatic factors (De Bruijn, 1973; Bokanga *et al.*, 1994), as evidenced by the correlation between drought periods in Africa and increased cassava flour toxicity. Further, these periods correspond with an increased incidence of cyanide poisoning and outbreaks of Konzo within communities reliant on cassava (Cliff, 1994; Cardoso *et al.*, 1999; Oluwole, 2015).

Climate change projections for cassava-growing regions in Africa include mean surface air temperature increases of 3–4 °C, with seasonal increases of up to 7 °C by 2099 (Collins *et al.*, 2013; Niang *et al.*, 2014). In general, models forecast an increase in aridity over most of Africa (Dai, 2011), and decreases in soil moisture and increased risk of agricultural drought in southern Africa (Collins *et al.*, 2013; Niang *et al.*, 2014). These forecasts highlight the need to investigate the effects of both drought and temperature on cassava.

A number of studies have investigated the impact of drought on cassava yield and productivity (e.g. Connor *et al.*, 1981; Keating *et al.*, 1982; El-Sharkawy & Cock, 1987; Baker *et al.*, 1989; El-Sharkawy *et al.*, 1992a; Bokanga *et al.*, 1994; El-Sharkawy & Cadavid, 2002; El-Sharkawy, 2006; Bakayoko *et al.*, 2009), but relatively few of these largely field-based studies have also measured cyanogenic capacity. Typically, the cyanide potential of tubers (CN_p; the maximum amount of cyanide released from all endogenous cyanogenic glucosides) is higher when soil moisture is low (e.g. De Bruijn, 1973; Santisopasri *et al.*, 2001; Okogbenin *et al.*, 2003). Even fewer studies report CN_p of leaves, even though leaves are an important animal feed or protein supplement for humans (Gomez *et al.*, 1985; Ngudi *et al.*, 2003). A controlled glasshouse study investigated the effects of drought on the growth and chemistry of cassava during the early stages of tuber development, and found significant increases in tuber and leaf CN_p in drought plants after 14–28 days of water deficit (Vandegeer *et al.*, 2013). This increase in toxicity was reported in plants grown at air temperatures at the lower end of the range at which cassava grows (18.8/16.9 °C mean day/night temperatures). Presumably higher temperatures would exacerbate the effects of drought.

Cassava is highly plastic in its growth response to air temperature. Studies of the effects of temperature on cassava have tended to focus either on low temperature limitations to yield and effects on biomass allocation (Cock & Rosas, 1975; Irikura *et al.*, 1979; Manrique, 1992; Fermont *et al.*, 2009), or on its photosynthetic capacity and growth under higher temperatures (e.g. Cock *et al.*, 1979; Edwards *et al.*, 1990; El-Sharkawy & Cock, 1990; El-Sharkawy *et al.*, 1992b). These studies generally report reduced yields and growth at temperatures less than 17 °C (Cock & Rosas, 1975; El-Sharkawy

et al., 1992b), and broad photosynthetic temperature optima at leaf temperatures between 25 and 40 °C (Mahon *et al.*, 1977; El-Sharkawy *et al.*, 1984, 1992b; El-Sharkawy & Cock, 1990). In addition, greater stimulation of cassava yield by elevated atmospheric CO₂ concentrations (700 ppm) was found at higher growth temperatures in a glasshouse pot trial under well-watered conditions (Imai *et al.*, 1984), further highlighting the importance of interactions between temperature and other factors in the response of cassava to changing climates.

Despite the often substantial effects of temperature on growth, photosynthesis and biomass partitioning of cassava, to our knowledge no studies to date have studied the effect of temperature on the toxicity of cassava, nor investigated interactive effects of drought and temperature on growth, yield and nutritional value. Given the importance of cyanogenic capacity of this staple to human nutrition, the general importance of food quality to achieving food security (Jarvis *et al.*, 2012; Van Rijssen *et al.*, 2013), predictions of increased temperatures and aridity in Africa, and the projected use of cassava (Scott *et al.*, 2000), understanding environmental effects on cyanogenic glucoside content is crucial.

Here, we present results of a controlled glasshouse experiment in which we examined the effects of both temperature and drought on the toxicity (for the first time), growth and biomass allocation of cassava. Specifically, we sought to address the following questions: (1) Under drought conditions, does higher temperature exacerbate the effects of water deficit on cyanogenic glucoside concentration? and (2) under well-watered conditions, is enhanced growth under optimal (typically high) temperatures at the expense of investment in defence? Results are discussed in the context of food security – both in terms of the amount and quality of food – and climate change forecasts for cassava-growing regions.

Materials and methods

Plant material and growing conditions

Forty-three cassava plants (*Manihot esculenta* Crantz cv. MCol 1468) were propagated clonally (as ~50 mm long cuttings) in sand, from a single parent plant. Thirty-eight cuttings had sprouted 58 days after cultivation and were transferred to individual 250-mm-diameter, plastic-free draining pots, containing 9 kg of a 1 : 4 (w/w) soil : sand mix. The soil : sand mix comprised washed river sand and soil from 0 to 150 mm depth in Jock Marshall Reserve, Monash University, Clayton, Victoria (37°54' S, 145°8' E), sieved to < 2 mm. This mixture, referred to as 'soil' hereafter, had low endogenous levels of plant-available (Colwell) phosphorus (3 mg kg⁻¹), mineral nitrogen (3 mg kg⁻¹ as the sum of NO₃⁻-N and NH₄⁺-N),

0.02% total nitrogen, 0.35% total carbon, a pH of 6.2, and high mycorrhizal fungal inoculum potential (R. E. Miller, unpub data). A 10-mm layer of polystyrene beads was placed on the soil surface to reduce evaporation.

For the first 85 days after planting (DAP), plants were watered as required. From 86 DAP, plants were watered every second day to field capacity (FC) (Khan *et al.*, 2003) with a modified Hoagland's solution containing 5 mM nitrogen (supplied as NO_3^- and NH_4^+ in the ratio 5 : 1), which a preliminary experiment showed to be optimum for growth (data not shown). From 114 DAP, plants were watered alternately with water and the nutrient solution; from 128 DAP, plants were watered with nutrient solution every third watering; and after 144 DAP, at which point the drought treatment was applied, only water was applied to avoid differences in nutrient supply to drought and well-watered plants.

For the first 80 DAP, plants were grown in a glasshouse with ambient temperature (mean day/night cycle of 22/18 °C) and natural light (22nd February to 12th May, 2011, Melbourne, Australia). At 81 DAP, plants were randomly allocated to one of two glasshouses and provided with supplemental lighting with a 16-/8-hour day/night photoperiod (MK-1 Just-a-shade, Ablite, Melbourne, Australia), and a photosynthetically active radiation (PAR) of $400 \pm 100 \mu\text{mol quanta m}^{-2} \text{s}^{-1}$ (LI-1400 Light Meter; Li-Cor Environmental, Nebraska, USA). Temperatures were maintained at day/night mean temperatures of 25/20 °C, within cassava's optimal range (El-Sharkawy, 2004). These conditions were maintained until 132 DAP when temperature treatments were imposed.

Treatments

To quantify the combined effects of temperature and water supply on cassava, plants were randomly assigned to treatments in a 2 × 2 factorial design. From 133 DAP, two temperature treatments were imposed; in one glasshouse, day/night mean temperatures were increased to 34/28 °C ($n = 19$ plants), referred to as 'high'; and in the other, 23/23 °C ($n = 19$), referred to as 'low'. The mean maximum day and minimum night temperatures in the high- and low-temperature glasshouses were 38 °C/25 °C and 26 °C/20 °C, respectively. The temperature in the high glasshouse was chosen based on IPCC (2013) projections of 7 °C warming in Africa by 2099 (Niang *et al.*, 2014), compared, for example, to present mean monthly temperatures of 29 °C in Mozambique (INAM, 2013, Vandeger *et al.*, 2013). Temperatures in the low glasshouse were selected to provide a large difference in growth temperatures between treatments, without diverging too far from realistic growth conditions for cassava. Plants were swapped two times between adjacent glasshouses during the treatment period to reduce potential glasshouse effects. From 144 DAP, watering regimes were applied (Fig. 1). Plants allocated to the well-watered ($n = 9$) treatment were watered to field capacity for the duration of the experiment. The drought treatment ($n = 10$) was imposed by withholding water until a soil moisture content of 25% field capacity was reached, following Vandeger *et al.* (2013). Plants in the drought treatment were maintained at 25% field capacity for the remainder

of the experiment. In summary, temperature treatments began four and a half months after striking cuttings and lasted 6 weeks, while the drought treatment began 5 months after striking and lasted 4 weeks.

Harvest and sampling

All plants were destructively harvested 176 DAP. As cassava was propagated clonally, the biomass of the original cutting was excluded from biomass measurements to account for variation in original cutting size (Vandeger *et al.*, 2013). Leaf area was determined using a leaf area meter (LI-3000 Portable Area Meter and LI-3050A Belt Conveyor; Li-Cor Inc., Lincoln, NE, USA). Above-ground biomass (stems, leaves) was dried at 60 °C for 7 days, for dry weight determination and chemical analysis.

Roots were washed free from the soil with water and separated into tubers (roots >5 mm diameter) and fine roots. A subsample of fine roots was stored in 70% ethanol for the determination of mycorrhizal colonization of roots (Appendix S1). Subsamples of the inner tuber flesh (parenchyma) were taken from the middle, longitudinally. Two samples (ca. 2 g fresh weight) were taken from the centre of these slices, avoiding the tuber peel (cortex). One sample was used for the determination of cyanogenic glucosides and the other for nitrogen and carbon analyses. Sections of tuber peel were also sampled from the middle of each tuber for the same analyses. All fine and coarse root material was dried at 60 °C for 7 days for




Fig. 1 Soil moisture content of cassava grown in two temperature × watering treatments. Plants were grown in high- (mean 34 °C) or low-temperature (mean 23 °C) glasshouses for 6 weeks from 133 days after planting (DAP) (solid arrow), and under either well-watered (100% field capacity) or drought (25% field capacity) conditions for 4 weeks from 144 DAP (dashed arrow). Treatments were low temperature well-watered (○), low temperature drought (●), high temperature well-watered (▽) and high temperature drought (▼). Data are means ± SE of $n = 9$ –10.

the determination of dry weight and per cent dry matter of all tubers. Harvest index was calculated by dividing the total tuberous root dry weight by the total plant dry weight.

Analytical methods

Cyanogenic glucoside concentrations. At 176 DAP, prior to destructive harvesting two leaf discs of 5 mm diameter were sampled from the middle of the centre lobe (avoiding the midrib) of the third fully expanded leaf of each plant, for analysis of cyanogenic glucosides. Cyanogenic glucoside concentrations were also determined for the two largest tubers from each plant. To avoid potential confounding effects of intratuber variation (Bradbury *et al.*, 1991), cyanide content of the middle section, longitudinally, of each tuber was used for analysis. Cyanogenic glucosides were measured as CNp, that is the total amount of cyanide (CN) evolved from fresh leaf or tuber tissue, according to Vandegeer *et al.* (2013). Cyanide captured in a well of 1 M NaOH was quantified using a colorimetric assay. Absorbance was measured at 595 nm with NaCN as the standard. Leaf discs and tuber samples were rinsed and dried in a 60 °C oven for 48 h to enable determination of mass-based cyanide concentrations.

To ensure there was no potential epigenetic effect of tissue age CNp (Jørgensen *et al.*, 2005), foliar and tuber flesh CNp were compared in plants derived from cuttings taken from different parts of the parent plant. CNp was not dependent on the position from which the cutting was obtained (data not shown). Further, both within and across all treatments, no significant size effect on tuber CNp was found; thus, differences in tuber CNp were not a consequence of any differences in tuber developmental stage (data not shown).

Elemental analyses. Dried leaf, tuber flesh and peel subsamples were ground to a fine powder in a cooled IKA Labortechnik A10 Analytical Mill (Janke & Kunkel, Stanfen, Germany). For each tissue, the concentration of nitrogen (N%) was determined for 4–7 mg dwt of tissue by dry combustion using an elemental analyser (Vario Micro Cube CHNS Analyser; Elementar Australia Pty Ltd., Sydney, Australia). To estimate the degree of water stress (Farquhar *et al.*, 1989), 3 mg dwt of tuber flesh was analysed for carbon isotopes ($\delta^{13}\text{C}_{\text{‰}}$) using an ANCA GSL2 elemental analyser coupled with a Hydra 20–22 isotope ratio mass spectrometer (Sercon Ltd., Crewe, Cheshire, UK) with a precision of 0.1‰. As cassava drops leaves in response to drought (e.g. Vandegeer *et al.*, 2013), we measured tuber $\delta^{13}\text{C}$ rather than foliar $\delta^{13}\text{C}$ to provide integrated measure of plant water use efficiency (WUE) over the entire time of tuber development (Farquhar *et al.*, 1989; Jefferies & Mackerron, 1997).

Foliar chlorophyll fluorescence, foliar chlorophyll concentration and root arbuscular mycorrhizal colonization were also measured. Method information for these analyses is in Appendix S1 (Supporting Information).

Statistical analysis

Data were analysed using two-factor general linear models (GLMs) in JMP v.9 (SAS Institute Inc., Cary, NC, USA, 2010)

and IBM SPSS Statistics for Windows 21.0 (IBM Corp, Armonk, NY, USA, 2012) statistical software. Where necessary, data were transformed to satisfy the assumptions of GLMs. Tukey's HSD tests were used post hoc to compare means at $P < 0.05$ where no significant interaction between water regime (*W*) and temperature (*T*) was detected. Where a significant $T \times W$ interaction was detected, simple main effects tests were conducted within temperature treatments to compare drought and well-watered treatments.

Results

Plant growth and physiology

At the final harvest, both temperature and drought treatments had influenced the growth of plants, as indicated by a significant interaction for total plant biomass ($F_{1,34} = 6.330$, $P = 0.017$; Table 1); the reduction in biomass with drought was greater at high temperature than low temperature. Total biomass was greatest in the high-temperature well-watered treatment (42 ± 3.5 g dwt; mean \pm 1SE), twofold greater than plants from both drought treatments and 1.5 times greater than plants from the low-temperature well-watered treatment. Irrespective of water treatment, plants grown at high temperature had a 1.5-fold increase in height ($F_{1,34} = 50.82$, $P < 0.001$; Fig. 2) and above-ground biomass ($F_{1,34} = 17.916$, $P < 0.001$; Fig. 3), compared with plants grown at low temperatures. There was no difference in the height of plants assigned to different temperature treatments prior to the application of temperature treatments; however, within 1 week of changing glasshouse temperatures, the growth rate of plants in the high-temperature glasshouse (20 mm day^{-1}) was twice that of plants in the low-temperature glasshouse (10 mm day^{-1}), a difference which persisted until harvest (Fig. 2).

Plants produced significantly more leaf biomass ($F_{1,34} = 7.720$, $P = 0.009$) and increased leaf area ($F_{1,34} = 4.480$, $P = 0.042$; Table 1) in the high-temperature treatment, and well-watered treatment ($F_{1,34} = 10.977$, $P = 0.002$; and $F_{1,34} = 5.577$, $P = 0.024$, respectively) with no interactive effects. Specific leaf area of leaves retained on the plant at harvest ranged from 250 to 290 $\text{cm}^2 \text{ g}^{-1}$ dwt, with no difference between treatments. Plants produced 15% more leaves (Table 1) in the high-temperature glasshouse ($F_{1,34} = 4.777$, $P = 0.036$); however, within 2 weeks of applying the drought treatment, drought-treated plants shed 40% more leaves ($F_{1,34} = 6.935$, $P = 0.013$; Table 1).

The total below-ground biomass of plants was influenced by the interactive effects of temperature and water treatments ($F_{1,34} = 9.461$, $P = 0.004$; Table 1, Fig. 3), with a greater magnitude reduction in root

Table 1 Mean (\pm SE) plant growth characteristics for cassava grown in low-temperature (mean 23 °C/23 °C, day/night) or high-temperature (mean 34 °C/28 °C) glasshouses, and either well-watered (100% field capacity, $n = 9$) or drought (25% field capacity, $n = 10$) treatments

Tissue/parameter	High temperature		Low temperature		GLM (<i>P</i>)		
	Well	Drought	Well	Drought	<i>T</i>	<i>W</i>	<i>T</i> × <i>W</i>
Whole plant							
Total biomass (g dwt)	41.9 \pm 3.5 ^a	19.4 \pm 1.9 ^b	28.4 \pm 4.6 ^A	21.5 \pm 2.0 ^A	ns	<0.001	0.017
Root : shoot	1.9 \pm 0.1 ^a	0.6 \pm 0.1 ^b	1.9 \pm 0.2 ^A	1.6 \pm 0.1 ^A	<0.001	<0.001	<0.001
Harvest index (%)	54.7 \pm 1.9 ^a	23.1 \pm 1.7 ^b	53.5 \pm 3.7 ^A	48.7 \pm 2.98 ^A	<0.001	<0.001	<0.001
Shoots							
Total number leaves*	27.0 \pm 1.4 ^a	27.6 \pm 1.4 ^a	23.9 \pm 1.7 ^b	24.0 \pm 1.6 ^b	0.036	ns	ns
Number leaves dropped	10.2 \pm 0.5 ^{ab}	14.3 \pm 1.4 ^a	8.7 \pm 1.0 ^b	11.5 \pm 1.8 ^{ab}	ns	0.013	ns
Number leaves retained	16.8 \pm 1.2 ^a	13.3 \pm 0.4 ^b	15.2 \pm 0.9 ^{ab}	12.5 \pm 0.6 ^b	ns	<0.001	ns
Leaf mass (g dwt)**	5.2 \pm 0.3 ^a	3.8 \pm 0.2 ^b	3.9 \pm 0.5 ^{ab}	3.2 \pm 0.3 ^b	0.009	0.002	ns
Leaf area (cm ²)**	1292 \pm 77.3 ^a	1005 \pm 69.8 ^{ab}	1026 \pm 119.7 ^{ab}	911 \pm 69.0 ^b	0.042	0.024	ns
Roots							
Tuber % dry matter	30.5 \pm 0.5 ^a	20.6 \pm 0.9 ^{ba}	28.1 \pm 1.6 ^A	27.6 \pm 0.6 ^A	0.029	<0.001	<0.001
Tuber biomass (g dwt)	23.3 \pm 2.6 ^a	4.6 \pm 0.7 ^b	16.4 \pm 3.2 ^A	10.7 \pm 1.4 ^A	ns	<0.001	0.004
Fine root mass (g dwt)	4.1 \pm 0.3 ^a	2.8 \pm 0.2 ^b	2.8 \pm 0.4 ^b	2.4 \pm 0.2 ^b	0.002	0.002	ns

Results (*P* values) of two-way general linear models (GLMs) of temperature (*T*) and water regime (*W*) are shown. Significant differences between means are indicated by superscript letters (Tukey's HSD; $P < 0.05$) across all treatments, or within temperature treatments when the *T* × *W* interaction was significant.

*Total number of leaves over course of experiment.

**Leaf mass and leaf area measured for leaves retained at harvest.




Fig. 2 Height (cm) of cassava plants subject to two temperature × watering treatments measured at intervals throughout the experiment until harvest 176 days after planting (DAP). Plants were grown in high-temperature (mean 34 °C) or low-temperature (mean 23 °C) glasshouses for 6 weeks from 133 DAP (solid arrow), and under either well-watered (100% field capacity) or drought (25% field capacity) conditions for 4 weeks from 144 DAP (dashed arrow). Treatments were low temperature well-watered (○), low temperature drought (●), high temperature well-watered (▽) and high temperature drought (▼). Data are means \pm SE of $n = 9$ –10. *indicates significant differences between temperature treatments at $P < 0.05$.

biomass at high temperature than low temperature. Specifically, and similar to above-ground biomass, total below-ground biomass was greatest in the high-temperature well-watered treatment (27.4 ± 2.7 g dwt; mean \pm 1SE), four times higher than plants in the high-temperature drought treatment. The same pattern was observed for tuber biomass ($F_{1,34} = 9.717$, $P = 0.004$; Table 1). The difference in tuber mass was not a consequence of differences in the number of tubers, which was similar across all treatments (overall mean \pm 1SE of 2.3 ± 0.15 tubers per plant). Significant water × temperature effects were detected for root:shoot ($F_{1,34} = 27.01$, $P < 0.001$) and harvest index ($F_{1,34} = 25.57$, $P < 0.001$; Table 1). Specifically, harvest index (23 ± 1.7 , mean \pm 1SE) and root:shoot (0.6 ± 0.05 , mean \pm 1SE) of plants from the high-temperature drought treatment were less than half all other treatments. Plants produced more fine roots (Table 1) in the high temperature ($F_{1,34} = 11.64$, $P = 0.002$) and well-watered treatments ($F_{1,34} = 10.76$, $P = 0.002$), with no interactive effects. Under well-watered conditions, increases in above- and below-ground biomass with high temperature were proportional, and temperature therefore had no detectable effect on biomass partitioning (harvest index and root:shoot; Table 1). Across all treatments, mycorrhizal fungi colonized over 95% of fine root length, with no difference between treatments detected (data not shown).




Fig. 3 Above- and below-ground biomass of cassava plants grown in two temperature \times watering treatments. Treatments were high (HT, mean 34 °C; white bars) or low temperature (LT, mean 23 °C; light grey bars) imposed for 6 weeks from 133 days after planting (DAP), and well-watered (W, 100% field capacity; open bars) or drought (D, 25% field capacity; hatched bars) imposed for 4 weeks from 144 days after planting (DAP). Fine root biomass (dark grey) and tuber biomass are shown. Data are means \pm SE of $n = 9$ –10. Results of two-way GLMs are shown for total shoot and total root biomass; letters indicate significant differences at $P < 0.05$. Where the $T \times W$ interaction was significant, letters indicate significant differences between drought and well-watered treatments at each temperature.

Changes in mean per cent dry matter of tubers reflected a significant interaction between drought and temperature treatments ($F_{1,34} = 21.537$, $P < 0.001$; Table 1). At low temperature, tuber dry matter (%) was similar between drought and well-watered plants, whereas at high temperature, drought effected a significant reduction in tuber dry matter (%) from a maximum of $30.5 \pm 0.5\%$ (mean \pm 1 SE) under well-watered conditions to a minimum of $20.6 \pm 0.9\%$ (mean \pm 1 SE) under drought (Table 1).

The photosynthetic efficiency (F_v/F_m) of all plants one week before harvest was 0.76 ± 0.1 (mean \pm 1 SE) with no difference between drought or temperature treatments (data not shown). Total chlorophyll concentration in the third fully expanded leaf of each plant was 1.3-fold higher in drought plants and did not differ with temperature ($F_{1,34} = 14.15$, $P < 0.001$; Appendix S1).

Plant chemical composition

$\delta^{13}\text{C}$ was determined on tubers that were likely initiated under equivalent conditions (approx. 3 months after planting; Alves, 2002) but developed for 6 weeks under treatment conditions. Plants had tuber $\delta^{13}\text{C}$ values ranging from -25.9‰ to -22‰ , with highest values in the low temperature ($F_{1,34} = 436.3$, $P < 0.001$) and drought treatments ($F_{1,34} = 131.4$, $P < 0.001$; Table 2), with no interactive effects. There was no significant correlation between tuber $\delta^{13}\text{C}$ and tuber cyanide concentration within or across all treatments (data not shown).

Table 2 Mean (\pm SE) chemical composition of tissues from cassava grown in low temperature (mean 23 °C/23 °C, day/night) or high temperature (mean 34 °C/28 °C) glasshouses, and either well-watered (100% field capacity, $n = 9$) or drought (25% field capacity, $n = 10$) treatments

Tissue type/parameter	High temperature		Low temperature		GLM (P)		
	Well	Drought	Well	Drought	T	W	T \times W
Leaves							
CN-N/N (%)*	3.9 \pm 0.5	4.3 \pm 0.5	3.2 \pm 0.4	4.9 \pm 0.9	ns	ns	ns
C : N	17.6 \pm 0.2 ^a	13.3 \pm 0.2 ^b	16.8 \pm 0.5 ^A	14.7 \pm 0.4 ^B	ns	<0.001	0.002
Tuber							
CN-N/N (%)*	1.5 \pm 0.2 ^a	5.5 \pm 1.0 ^b	7.1 \pm 1.8 ^A	6.9 \pm 1.3 ^A	0.002	0.020	0.007
C : N	275.7 \pm 16.2 ^a	125.2 \pm 10.4 ^b	197.4 \pm 24.2 ^A	148.3 \pm 7.6 ^A	ns	<0.001	0.002
$\delta^{13}\text{C}$ (‰)	-25.9 \pm 0.14 ^a	-24.6 \pm 0.11 ^b	-23.5 \pm 0.11 ^c	-22.0 \pm 0.09 ^d	<0.001	<0.001	ns
Tuber peel							
CNp (mg g ⁻¹ dwt)	0.77 \pm 0.20 ^a	1.63 \pm 0.25 ^{ab}	1.77 \pm 0.60 ^{ab}	3.19 \pm 0.52 ^b	0.005	0.012	ns
Nitrogen (mg g ⁻¹ dwt)	3.14 \pm 0.22 ^a	8.26 \pm 0.74 ^b	5.94 \pm 0.64 ^A	6.40 \pm 0.43 ^A	ns	<0.001	<0.001
CN-N/N (%)*	13.1 \pm 5.3 ^a	11.3 \pm 2.5 ^a	25.9 \pm 8.3 ^b	35.8 \pm 3.7 ^b	0.001	ns	ns
C : N	135.7 \pm 7.0 ^a	54.8 \pm 4.5 ^b	75.9 \pm 7.2 ^A	66.3 \pm 4.2 ^A	0.0003	<0.001	<0.001

Results (P values) of two-way general linear models (GLMs) of temperature (T) and water regime (W) are shown. Significant differences between means are indicated by superscript letters (Tukey's HSD; $P < 0.05$) across all treatments, or within temperature treatments when the $T \times W$ interaction was significant.

*CN-N/N (%) is the proportion of total N that is present as CN in each tissue type.

The nitrogen (N) concentration of leaves (28.35–37.17 mg g⁻¹ dwt; Fig. 4a) was higher than that of tuber peel (3.14–8.26 mg g⁻¹ dwt; Table 2) and tuber flesh (1.52–3.51 mg g⁻¹ dwt; Fig. 4c), across all treatments. There was a significant interactive effect of temperature and water treatments on N concentration in leaves ($F_{1,33} = 8.286$, $P = 0.006$), tuber flesh ($F_{1,34} = 7.538$, $P = 0.010$) and tuber peel ($F_{1,21} = 22.03$, $P < 0.001$). Nitrogen concentrations in all tissues were highest in the high-temperature drought treatment and lowest in the high-temperature well-watered treatment, with a trend towards higher N in tissues of drought plants. The magnitude of the drought effect differed between temperature treatments such that at high-temperature tuber flesh and peel N increased 2.3-fold and

2.6-fold with drought, respectively, but at low temperature no differences with drought were detected (Table 2, Fig. 4c). Changes in tissue carbon-to-nitrogen ratios (C : N) reflected the changes in N concentrations and not in C, with a significant reduction in C : N under drought in all tissues that was of greater magnitude under high temperature than low temperature (Table 2).

A significant effect of temperature alone on the concentration of cyanogenic glucosides (CNp) was only observed in well-watered plants. Across all treatments, CNp was highest in tuber peel (0.77–3.19 mg g⁻¹ dwt; Table 2) and leaves (1.78–3.21 mg g⁻¹ dwt; Fig. 4b), and lowest in the tuber flesh (0.05–0.35 mg g⁻¹ dwt; Fig. 4d). For all tissues, a significant main effect of




Fig. 4 Nitrogen concentration (a, c) and cyanide potential (CNp) (b, d) of cassava leaves and tubers from plants grown in two temperature × watering treatments. Plants were grown in high- (mean 34 °C; white bars) or low-temperature (mean 23 °C; light grey bars) glasshouses for 6 weeks from 133 days after planting (DAP), and in well-watered (100% field capacity; open bar) or drought (25% field capacity; hatched) conditions for 4 weeks from 144 DAP. Data are means ± SE of $n = 9-10$. Results of two-way GLMs are shown; different letters indicate significant differences at $P < 0.05$. Where the $T \times W$ interaction was significant, letters indicate significant differences between drought and well-watered treatments at each temperature.

watering regime was detected, with significant increases in CNp under drought (Table 2, Fig. 4). Drought effected an increase in leaf cyanide concentration, irrespective of temperature ($F_{1,34} = 8.778$, $P = 0.0061$), with 1.8-fold and 1.5-fold increases in leaf CNp in the low- and high-temperature treatments, respectively (Fig. 4b). No significant main effect of temperature on leaf CNp was detected. By contrast, significant differences in CNp with temperature were detected in tuber tissues. In tuber flesh, the magnitude of the drought effect differed significantly between temperature treatments, with 6.7-fold greater CNp in tubers from drought plants at high temperature, whereas at low temperature, the trend towards increased CNp with drought was not significant (Fig. 4d). Significant main effects of temperature ($F_{1,34} = 8.963$, $P = 0.005$) and watering treatment ($F_{1,34} = 7.136$, $P = 0.012$) were found on tuber peel CNp, which was greater under drought and at low temperature (Table 2). Pooling watering treatments, tuber peel CNp of low-temperature-grown plants was double that of high-temperature-grown plants.

The proportion of foliar N allocated to cyanogenic glucosides (CN-N/N%) was similar across all treatments (mean 4.1%), despite significant differences in CNp with drought (Table 2). By contrast, significant effects of temperature and drought on CN-N/N% were detected in tuber tissues. Specifically, at high temperature, a significantly greater proportion of N was allocated to CN under drought, whereas at low temperature CN-N/N% was similar between watering treatments. There was a significant main effect of temperature on tuber peel CN-N/N%, with on average higher CN-N/N% at low temperature (mean 30.9%) than high temperature (mean 12.2%; Table 2).

Discussion

We report here, for the first time, the effects of temperature, and temperature combined with drought on growth, biomass partitioning and nitrogen allocation to cyanogenic glucosides in cassava leaves and tubers under controlled conditions. Increases in tuber toxicity with drought and at lower growth temperatures point to potential trade-offs between growth and secondary metabolism in resource allocation. Greater complexity arises in the different responses of cyanogenesis in above- and below-ground tissues to drought and temperature. Findings are consistent with models that predict cassava to be adaptable and resilient to forecast temperature increases (Jarvis *et al.*, 2012), but show that irrespective of growth temperature, drought effects on tuber yield and toxicity are the greater threat to future food security.

Temperature effects on growth of well-watered plants

The high- (34 °C) and low-temperature (23 °C) regimes used here represent the low and high ends of the temperature range where cassava is cultivated (El-Sharkawy & Cock, 1990; El-Sharkawy *et al.*, 1992b). In accord with El-Sharkawy *et al.* (1992b), who found the optimum temperature range for cassava photosynthesis is 30–40 °C, and Mahon *et al.* (1977) who found higher photosynthetic and growth rates at 29/24 °C compared to 24/19 °C, the high-temperature treatment had significant and rapid effects on plant growth under well-watered conditions, clearly indicating that the low-temperature treatment was suboptimal for growth of this cultivar. Leaf growth in cassava is known to decrease at lower temperatures (Irikura *et al.*, 1979). We found no effect of temperature on biomass partitioning under well-watered conditions (Table 1); no consistent effect of temperature on biomass partitioning is evident (e.g. Mahon *et al.*, 1976; Keating *et al.*, 1982).

Impact of temperature on drought responses – yield, biomass and physiology

Above-ground, the physical responses to drought observed here are similar to previous studies with smaller, fewer leaves, at both growth temperatures. While drought alone drove changes in leaf loss and retention, both temperature and drought affected leaf area and biomass (Table 1). Leaf formation and growth in cassava are known to be highly sensitive to even small decreases in soil moisture (Connor & Cock, 1981; Connor *et al.*, 1981; Baker *et al.*, 1989; De Tafur *et al.*, 1997; Okogbenin *et al.*, 2003; Alves & Setter, 2004; Vandegeer *et al.*, 2013). Rapid closure of stomata, combined with leaf abscission in response to more prolonged water deficit, enable cassava to retain photosynthetically active, turgid leaves (El-Sharkawy & Cock, 1984; Palta, 1984; Alves, 2002; Turyagyenda *et al.*, 2013; Vandegeer *et al.*, 2013). Consistent with this, there was no reduction in photosynthetic efficiency (F_v/F_m) of leaves retained under drought (Calatayud *et al.*, 2002; see also Vandegeer *et al.*, 2013; but see Zhao *et al.*, 2015). By contrast, changes in tuber $\delta^{13}\text{C}$ across all treatments reflect the physiological effects of both temperature and drought treatments (Table 1). Consistent with high-temperature stimulation of photosynthesis and growth (discussed above), lower tuber $\delta^{13}\text{C}$ under well-watered and high-temperature conditions is indicative of higher intercellular CO_2 concentration (C_i) and stomatal conductance (g_s) (Farquhar *et al.*, 1989). In contrast, the less negative $\delta^{13}\text{C}$ of tubers of plants in the drought treatment indicates that the plants experienced some stress.

Tuber yield declines in response to drought are generally considered a consequence of reduction in canopy area and assimilate production (Connor & Cock, 1981; Baker *et al.*, 1989; De Tafur *et al.*, 1997; Setter & Fregene, 2007). Here, the relative yield decline with drought was greater at high temperature (80%) than at low temperature (35%, ns), but this was not explained by differences in leaf area, leaf biomass and leaf loss, which were largely similar at high and low temperatures. Only 24% of total biomass was in the tubers in the high-temperature drought plants – approximately half that in all other treatments which ranged from 50 to 58%. While some studies similarly report a greater relative decrease in tuber yield (93%) compared to shoot biomass (59%) with water stress (e.g. Aina *et al.*, 2007), others report a greater relative decrease in shoot biomass (e.g. El-Sharkawy, 2007; 18% and 57% in tubers and shoots, respectively, in one cultivar). Our data highlight the importance of growth temperature in affecting drought impacts on biomass allocation within a cultivar.

The reduction in tuber yield of 80% at high temperature here is within the range of yield reductions reported elsewhere with drought under both natural and controlled field conditions. For example, mean percentage declines in tuber biomass of 82–96% were found for nine cultivars of cassava when a water deficit (25% FC) was imposed early (28 DAP) and sustained until harvest between 3 and 6 months later (Aina *et al.*, 2007). Importantly, our data demonstrate that growth temperature, as well as timing and duration of water deficit, is important in determining the effect of drought on tuber yield (e.g. see Alves, 2002). While there have been no prior studies of temperature × drought effects on cassava, there is some indication that yield reductions in response to drought are more substantial under warmer temperatures. The yield reductions of 82–96% reported by Aina *et al.* (2007), for example, were at a mean maximum temperature of 32 °C, whereas at a mean growth temperature of 23 °C tuber yield only declined 0–25%, albeit using different cultivars (El-Sharkawy *et al.*, 1992a). In the latter study, similar to findings here, the absence of substantial yield declines (0–25%) with drought at 23 °C occurred despite drought plants showing other effects of water deficit including lower photosynthetic rates and reduced leaf area and shoot biomass at that temperature.

Temperature and drought effects on plant chemistry and nutritional value

Global temperatures are rising, but there are relatively few papers on the likely impact on cassava yield (e.g. Lobell *et al.*, 2008; Knox *et al.*, 2012), and none to our knowledge on the direct effects of temperature on

cyanogenic glucosides. Knowledge of temperature effects on cyanogenic glucosides in general is limited to a few studies on clover that has a low optimum growth temperature (e.g. Stochmal & Oleszek, 1997; Hayden & Parker, 2002). In the present study, enhanced growth of cassava at higher temperature under well-watered conditions was associated with a significant reduction in tuber CNp and in the proportion of N allocated to CN, but there was no change in foliar chemistry or N allocation (Fig. 4; Table 2), confirming field studies showing that foliar chemistry is not a suitable proxy for estimating tuber toxicity (Bokanga *et al.*, 1994; Jørgensen *et al.*, 2005; Burns *et al.*, 2012b). This result also supports the assertion that environmentally driven changes in foliar defence metabolites cannot be assumed to be representative of the whole plant, an important consideration for root food crops (Parker *et al.*, 2012; Miller *et al.*, 2014). The lower tuber CNp at high temperature may indicate a reallocation of N away from defence under conditions that stimulate growth (Herms & Mattson, 1992; Neilson *et al.*, 2013). A reallocation of resources is plausible, given that differences in tuber CNp with temperature under well-watered conditions could not be explained either by changes in biomass (i.e. tissue dilution) or nitrogen concentration.

Trade-offs between growth and defence are more likely to be evident under resource limitation (Coley *et al.*, 1985), as was the case here, where plants were N deficient (<4% leaf N) at the time of harvest (Reuter & Robinson, 1997; Howeler, 2002). Tissue N is positively correlated with foliar cyanogenic glucoside concentrations in some species (e.g. *Eucalyptus cladocalyx*, Gleadow & Woodrow, 2000; *Sorghum bicolor*, Busk & Møller, 2002), but not in others (e.g. *Prunus turneriana*, Miller *et al.*, 2004; *Beilschmiedia collina*, Miller & Tuck, 2013). Surprisingly, few studies have addressed the relationship between N and CN in cassava, or have tested the tubers. Some studies indicate a relationship between foliar N and CNp in the shoot apex and leaves of cassava (Jørgensen *et al.*, 2005) whereas other field and glasshouse studies report no correlation between N and CNp in either tubers or foliage (Gleadow *et al.*, 2009; Burns *et al.*, 2012b). Here, unlike for CNp, we could largely account for the decrease in tuber N by the increase in tuber size. Thus, it appears that tuber CNp is regulated independently of tuber (or foliar) N concentrations. This is not surprising given that while some synthesis of cyanogenic glucosides occurs in the roots (Du *et al.*, 1995; McMahon *et al.*, 1995), cyanogenic glucosides in cassava are predominantly synthesized in the shoots and transported to the roots (Jørgensen *et al.*, 2005).

For a plant such as cassava, with a high optimum growth temperature, a major consequence of climate

change will arise from the interaction between rising temperatures and drought on yield and, importantly, the concomitant changes in toxicity and nutritional value. Reduction in tuber quality, as measured by per cent dry matter, and starch yield under water stress have previously been reported (e.g. Santisopasri *et al.*, 2001; El-Sharkawy, 2007; Bakayoko *et al.*, 2009), consistent with the significant reduction in tuber quality reported here under high temperature/drought conditions, from 31.5% to 20.5% dry matter (Table 1).

Of almost 30 papers on cassava describing controlled environmental and field-based studies in which drought responses were reported, only seven measured the effect on tuber cyanogenic glucosides (El-Sharkawy, 1993; Bokanga *et al.*, 1994; Santisopasri *et al.*, 2001; Okogbenin *et al.*, 2003; El-Sharkawy, 2006; Hular-Bograd *et al.*, 2011; Vandegeer *et al.*, 2013), and none report the interactive effects of temperature and drought. Four studies report mean relative increases in tuber toxicity ranging from 54 to 82% across 27 cultivars, in 12-month-old plants subject to drought varying in timing and duration (El-Sharkawy, 1993; Bokanga *et al.*, 1994; Okogbenin *et al.*, 2003; El-Sharkawy, 2006). Here, within a population of clones from a single cultivar, we demonstrate the importance of the interactive effects of temperature and drought, with greatest relative increase in tuber toxicity with drought (600%) found at high temperature, in part as a consequence of lower tuber CN_p under well-watered conditions. As was the case with differences between temperature treatments under well-watered conditions, tissue dilution of cyanogenic glucoside content did not account for differences between treatments (see also Bokanga *et al.*, 1994). Changes in tuber mass with drought did not account for changes in tuber N either, as tuber N concentrations increased, but tuber N content (per tuber) halved.

Increases in foliar cyanogenic glucoside concentrations with drought are known from other species (e.g. Gleadow & Woodrow, 2002), but very little is known about how temperature may affect that response. Whereas temperature × drought effects on tuber flesh (and peel) chemistry were more complex, drought alone drove changes in foliar CN_p, with similar increases (mean 62%) at both temperatures (Fig. 4; Table 2). Further, changes in foliar N and foliar CN_p were largely proportional across all treatments (Fig. 4); thus, increased foliar N, CN_p and chlorophyll in drought plants could be consistent with reclamation of constituents from abscising leaves (Aerts, 1996; Munné-Bosch & Alegre, 2004), a process which has also been hypothesized to contribute to increased tuber CN_p under drought (Vandegeer *et al.*, 2013).

It is likely that the changes in above- and below-ground cassava tissue chemistry in response to temperature × drought treatments here reflect a combination of factors, including changes in biomass, reclamation and reallocation of nutrients, trade-offs in N allocation, as well as more direct environmental effects on biosynthesis and transport of cyanogenic glucosides. Many of these processes and factors are not yet fully understood. For example, we require a greater understanding of factors affecting transport, remobilization and biosynthesis of cyanogenic glucosides (Møller, 2010; Neilson *et al.*, 2013), which may be independently regulated in roots and shoots (Blomstedt *et al.*, 2012; Miller *et al.*, 2014). Moreover, interpreting distribution and allocation of N to cyanogenic glucosides with respect to their defensive function may be further limited by the increasingly recognized nondefensive roles of these metabolites in storage and moderating stress (Neilson *et al.*, 2013; Selmar & Kleinwächter, 2013; Gleadow & Møller, 2014).

Implications for food security in a changing climate

Achieving food security is not only about increasing yields, but also ensuring that food is safe and of sufficient nutritive value. Cassava yields are expected to be positively impacted by a warming climate, within the range currently projected (Lobell *et al.*, 2008; Jarvis *et al.*, 2012). We found that tuber yields were increased and CN_p was decreased when plants were grown under higher temperatures with adequate water supply; however, the combined effects of higher temperature with drought stress had a negative effect on tuber yields and nutritive value. This plasticity highlights the importance of considering climate change as a multifactor phenomenon, and has important implications for those that currently rely on cassava as their main source of food and income (see Nhassico *et al.*, 2008; Muoki & Maziya-Dixon, 2010). Little is known about how drought and temperature might interact with the effects of increasing atmospheric [CO₂] on cassava growth and chemistry. Greater elevated [CO₂] (700 ppm) stimulation of yield was found at a higher growth temperature (33 °C/31 °C day/night) than lower growth temperature (28 °C/21 °C) under well-watered conditions in a glasshouse pot study, pointing towards potentially additive or synergistic effects of increasing temperature and [CO₂] on cassava growth and yield (CN_p was not measured; Imai *et al.*, 1984). However, inconsistent findings with respect to elevated [CO₂] effects on cassava growth and chemistry have been reported in glasshouse (pot) and field-based studies (e.g. Gleadow *et al.*, 2009; Rosenthal *et al.*, 2012). The only CO₂ enrichment study under field conditions was

in fertile soils in the absence of drought and found a substantial (104%) increase in tuber biomass, as well as enhanced WUE and lower foliar N concentrations under elevated [CO₂] (585 ppm; Rosenthal *et al.*, 2012). How the physiological responses of cassava to increasing [CO₂], temperature and drought combine to affect both yield and nutritional value (toxicity) requires attention.

In addition to the effects of increasing [CO₂], the combined effects of high temperature and drought investigated here also have implications for efforts seeking to expand production of cassava as a security crop in the face of a changing climate (e.g. Rufino *et al.*, 2013). The importance of this is emphasized by projections suggesting that we will see an almost doubling of cassava utilization in sub-Saharan Africa from the 1990s to 2020 (Scott *et al.*, 2000). Over the latter part of the 20th century, an increase in temperature (1–2 °C/50 years) and decrease in precipitation (up to 1 mm day⁻¹ per 50 years) have been observed in much of southern Africa, and over this century, climate models forecast an increase in aridity across Africa, and faster increases in temperature in Africa than the global average (Dai, 2011; Niang *et al.*, 2014). These projections, coupled with our findings, suggest that careful attention needs to be paid to the yield and nutritive responses of cassava in major cassava-growing regions. This will be especially important where subsistence farmers have sole reliance on cassava in times of environmental and social stress (Cliff, 1994; Burns *et al.*, 2010).

Acknowledgements

We gratefully acknowledge The Australian Government's aid agency, AusAID for funding this research through a grant to TRC and RMG. We also thank Rebecca Vandegeer and Natalie O'Donnell for useful discussions and technical advice, Stewart Crowley and Cécile de Fondaumiere for help monitoring plants and facilities during the experiment, and Dr Karen Little, Andrew Longmire, Dr Stephanie Watts-Williams, Phil Falcke and Adele Neale for assistance with plant harvest. REM was in part supported by a Margaret Clayton Fellowship, and TRC by an ARC Future Fellowship (FT120100463). We also thank the reviewers for their constructive contributions during the review process.

References

- Aerts R (1996) Nutrient resorption from senescing leaves of perennials: are there general patterns? *Journal of Ecology*, **84**, 597–608.
- Aina OO, Dixon AGO, Akinrinde EA (2007) Effect of soil moisture stress on growth and yield of cassava in Nigeria. *Pakistan Journal of Biological Sciences*, **10**, 3085–3090.
- Alves AA (2002) Cassava botany and physiology. In: *Cassava: Biology, Production and Utilization*, (eds Hillocks RJ and Thresh JM Bellotti AC) pp. 67–89. CABI Publishing, Wallingford, Oxon, UK.
- Alves AA, Setter TL (2004) Response of cassava leaf area expansion to water deficit: cell proliferation, cell expansion and delayed development. *Annals of Botany*, **94**, 605–613.
- Bakayoko S, Tschannen A, Nindjin C, Dao D, Girardin O, Assa A (2009) Impact of water stress on fresh tuber yield and dry matter content of cassava (*Manihot esculenta* Crantz) in Côte d'Ivoire. *African Journal of Agricultural Research*, **4**, 021–027.
- Baker GR, Fukai S, Wilson GL (1989) The response of cassava to water deficits at various stages of growth in the subtropics. *Australian Journal of Agricultural Research*, **40**, 517–528.
- Blomstedt C, Gleadow RM, O'Donnell N *et al.* (2012) A combined biochemical screen and TILLING approach identifies mutations in *Sorghum bicolor* L. Moench resulting in acyanogenic forage production. *Plant Biotechnology Journal*, **10**, 54–66.
- Bokanga M, Ekanayake IJ, Dixon AGO, Porto MCM (1994) Genotype-environment interactions for cyanogenic potential in cassava. *Acta Horticulturae*, **375**, 131–139.
- Bradbury JH, Egan SV, Lynch MJ (1991) Analysis of cyanide in cassava using acid-hydrolysis of cyanogenic glucosides. *Journal of the Science of Food and Agriculture*, **55**, 277–290.
- Burns AE, Gleadow RM, Cliff J, Zacarias AM, Cavagnaro TR (2010) Cassava: the drought, war and famine crop in a changing world. *Sustainability*, **2**, 3572–3607.
- Burns AE, Gleadow RM, Bradbury JH, Cliff J, Cavagnaro TR (2012a) Cyanogens in commercial food products of cassava. *Journal of Food Composition and Analysis*, **25**, 79–82.
- Burns AE, Gleadow RM, Zacarias AM, Cuambe CE, Miller RE, Cavagnaro TR (2012b) Variations in the chemical composition of Cassava (*Manihot esculenta* Crantz) leaves and roots as affected by genotypic and environmental variation. *Journal of Agricultural and Food Chemistry*, **60**, 4946–4956.
- Busk PK, Møller BL (2002) Dhurrin synthesis in *Sorghum* is regulated at the transcriptional level and induced by nitrogen fertilization in older plants. *Plant Physiology*, **129**, 1222–1231.
- Calatayud PA, Barón CH, Velásquez H, Arroyave JA, Lamaze T (2002) Wild *Manihot* species do not possess C4 photosynthesis. *Annals of Botany*, **89**, 125–127.
- Cardoso AP, Ernesto M, Cliff J, Bradbury JH (1999) High levels of total cyanogens in cassava flour related to drought in Mozambique. *Roots*, **6**, 4–6.
- Cliff J (1994) Cassava safety in times of war and drought in Mozambique. *Acta Horticulturae*, **375**, 372–378.
- Cock JH, Rosas S (1975) Ecophysiology of cassava. In: *Symposium on Ecophysiology of Tropical Crops*. pp. 1–14. Communications Division of CEPLAC, Ilheus-Itabuna, Brazil.
- Cock JH, Franklin D, Sandoval G, Juri P (1979) The ideal cassava plant for maximum yield. *Crop Science*, **19**, 271–279.
- Coley PD, Bryant JP, Chappin FS III (1985) Resource availability and plant antiherbivore defense. *Science*, **230**, 895–899.
- Collins M, Knutti R, Arblaster J *et al.* (2013) Long-term climate change: projections, commitments and irreversibility. In: *Climate Change 2013: The Physical Science Basis* (eds Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Stocker TF, Qin D, Plattner GK, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex B, Midgley BM]). Cambridge University Press, Cambridge, UK/New York.
- Conn EE (1981) Cyanogenic glycosides. In: *The Biochemistry of Plants* (ed. Conn EE) pp. 479–499. Academic Press Inc, New York.
- Connor DJ, Cock JH (1981) Response of cassava to water shortage. II. Canopy dynamics. *Field Crops Research*, **4**, 185–196.
- Connor DJ, Cock JH, Parra GE (1981) Response of cassava to water shortage. I. Growth and yield. *Field Crops Research*, **4**, 181–200.
- Dai A (2011) Drought under global warming: a review. *WIREs Climate Change*, **2**, 45–65.
- De Bruijn GH (1973) The cyanogenic character of cassava (*Manihot esculenta*). In: *Chronic Cassava Toxicity, Proceedings of an Interdisciplinary Workshop*, (eds Nestel BL and MacIntyre R) pp. 43–48 IDRC, Ottawa, Canada.
- De Tafur SM, El-Sharkawy MA, Cadavid LF (1997) Response of cassava (*Manihot esculenta* Crantz) to water stress and fertilization. *Photosynthetica*, **34**, 233–239.
- Du L, Bokanga M, Møller BL, Halkier BA (1995) The biosynthesis of cyanogenic glucosides in roots of cassava. *Phytochemistry*, **39**, 323–326.
- Edwards GE, Sheta E, Moore BD *et al.* (1990) Photosynthetic characteristics of cassava (*Manihot esculenta* Crantz), a C3 species with chlorenchymatous bundle sheath cells. *Plant and Cell Physiology*, **31**, 1199–1206.
- El-Sharkawy MA (2004) Cassava biology and physiology. *Plant Molecular Biology*, **56**, 481–501.
- El-Sharkawy MA (2006) International research on cassava photosynthesis, productivity, eco-physiology, and responses to environmental stresses in the tropics. *Photosynthetica*, **44**, 481–512.
- El-Sharkawy MA (2007) Physiological characteristics of cassava tolerance to prolonged drought in the tropics: Implications for breeding cultivars adapted to

- seasonally dry and semiarid environments. *Brazilian Journal of Plant Physiology*, **19**, 257–286.
- El-Sharkawy MA, Cadavid LF (2002) Response of cassava to prolonged water stress imposed at different stages of growth. *Experimental Agriculture*, **38**, 333–350.
- El-Sharkawy MA, Cock JH (1984) Water use efficiency of Cassava. I. Effects of air humidity and water stress on stomatal conductance and gas exchange. *Crop Science*, **24**, 497–502.
- El-Sharkawy MA, Cock JH (1987) Response of cassava to water stress. *Plant and Soil*, **100**, 345–360.
- El-Sharkawy MA, Cock JH (1990) Photosynthesis of cassava (*Manihot esculenta*). *Experimental Agriculture*, **26**, 325–340.
- El-Sharkawy MA (1993) Drought tolerance cassava for Africa, Asia and Latin America. *BioScience*, **43**, 441.
- El-Sharkawy MA, Cock JH, Held AA (1984) Photosynthetic responses of cassava cultivars (*Manihot esculenta* Crantz) from different habitats to temperature. *Photosynthesis Research*, **5**, 243–250.
- El-Sharkawy MA, Hernandez AD, Hershey C (1992a) Yield stability of cassava during prolonged mid-season water-stress. *Experimental Agriculture*, **28**, 165–174.
- El-Sharkawy MA, Tafur SMD, Cadavid LF (1992b) Potential photosynthesis of cassava as affected by growth conditions. *Crop Science*, **32**, 1336–1342.
- FAO & IFAD (2000) *The World Cassava Economy: Facts, Trends and Outlooks*. Food and Agriculture Organization of the United Nations and International Fund for Agricultural Development, Rome.
- Farquhar GD, Ehlering JR, Hubick KT (1989) Carbon isotope discrimination and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, **40**, 503–537.
- Fermont AM, Van AP, Giller KE (2008) Increasing land pressure in East Africa: the changing role of cassava and consequences for sustainability of farming systems. *Agriculture Ecosystems & Environment*, **128**, 239–250.
- Fermont AM, Van AP, Tittonell P, Van WM, Giller KE (2009) Closing the cassava yield gap: an analysis from smallholder farms in East Africa. *Field Crops Research*, **112**, 24–36.
- Gleadow RM, Møller BL (2014) Cyanogenic glycosides: Synthesis, physiology, and phenotypic plasticity. *Annual Review of Plant Biology*, **65**, 155–185.
- Gleadow RM, Woodrow IE (2000) Polymorphism in cyanogenic glycoside content and cyanogenic β -glucosidase activity in natural populations of *Eucalyptus cladocalyx*. *Australian Journal of Plant Physiology*, **27**, 693–699.
- Gleadow RM, Woodrow IE (2002) Defense chemistry of cyanogenic *Eucalyptus cladocalyx* seedlings is affected by water supply. *Tree Physiology*, **22**, 939–945.
- Gleadow RM, Evans JR, McCaffery S, Cavagnaro TR (2009) Growth and nutritive value of cassava (*Manihot esculenta* Crantz) are reduced when grown in elevated CO₂. *Plant Biology*, **11**(Suppl. 1), 76–82.
- Gomez G, Valdivieso M, Noma AT (1985) The influence of cultivar and plant age on the chemical composition of field-grown cassava leaves and roots. *Qualitas Plantarum-Plant Foods for Human Nutrition*, **35**, 109–119.
- Hayden KJ, Parker IM (2002) Plasticity in cyanogenesis of *Trifolium repens* L.: inducibility, fitness costs and variable expression. *Evolutionary Ecology Research*, **4**, 155–168.
- Herms DA, Mattson WJ (1992) The dilemma of plants: to grow or defend. *The Quarterly Review of Biology*, **67**, 283–335.
- Howler RH (2002) Cassava mineral nutrition and fertilization. In: *Cassava: Biology, Production and Utilization* (eds Hillocks RJ, Thresh JM, Bellotti AC), pp. 115–147. CABI Publishing, Wallingford, Oxon, UK.
- Hular-Bograd J, Sarobol E, Rojanaridpiched C, Sriroth K (2011) Effect of supplemental irrigation on reducing cyanide content of cassava variety Kasetsart 50. *Kasetsart Journal - Natural Science*, **45**, 985–994.
- Imai K, Coleman DF, Yanagisawa T (1984) Elevated atmospheric partial pressure of carbon-dioxide and dry matter production of cassava (*Manihot esculenta* Crantz). *Japanese Journal of Crop Science*, **53**, 479–485.
- INAM (2013) *Climate Information: Nampula*. Available at: www.inam.gov.mz (12 August 2013).
- Irikura Y, Cock JH, Kawano K (1979) The physiological basis of genotype-temperature interactions in cassava. *Field Crops Research*, **2**, 227–239.
- Jarvis A, Ramirez-Villegas J, Herrera Campo BV, Navarro-Racines C (2012) Is cassava the answer to African climate change adaptation? *Tropical Plant Biology*, **5**, 9–29.
- Jefferies RA, Mackerron DKL (1997) Carbon isotope discrimination in irrigated and droughted potato (*Solanum tuberosum* L.). *Plant, Cell and Environment*, **20**, 124–130.
- Jørgensen K, Bak S, Busk PK, Sørensen C, Olsen CE, Puonti-Kaerlas J, Møller BL (2005) Cassava plants with a depleted cyanogenic glucoside content in leaves and tubers. Distribution of cyanogenic glucosides, their site of synthesis and transport, and blockage of the biosynthesis by RNA interference technology. *Plant Physiology*, **139**, 363–374.
- Keating BA, Evenson JP, Fukai S (1982) Environmental effects on growth and development of cassava (*Manihot esculenta* Crantz). 3. Assimilate distribution and storage organ yield. *Field Crops Research*, **5**, 293–303.
- Khan HR, McDonald GK, Rengel Z (2003) Zn fertilization improves water use efficiency, grain yield and seed Zn content in chickpea. *Plant and Soil*, **249**, 389–400.
- Knox J, Hess T, Daccache A, Wheeler T (2012) Climate change impacts on crop productivity in Africa and South Asia. *Environmental Research Letters*, **7**, 034032.
- Lobell DB, Burke MB, Tebaldi C, Mastrandrea MD, Falcon WP, Naylor RL (2008) Prioritizing climate change adaptation needs for food security in 2030. *Science*, **319**, 607–610.
- Mahon JD, Lowe SB, Hunt LA (1976) Photosynthesis and assimilate distribution in relation to yield of cassava grown in controlled environments. *Canadian Journal of Botany*, **54**, 1322–1331.
- Mahon JD, Lowe SB, Hunt LA, Thiagarajah M (1977) Environmental effects on photosynthesis and transpiration in attached leaves of cassava (*Manihot esculenta* Crantz). *Photosynthetica*, **11**, 121–130.
- Manrique LA (1992) Growth and yield ability of cassava grown at 3 elevations in Hawaii. *Communications in Soil Science and Plant Analysis*, **23**, 129–141.
- McMahon JM, White WLB, Sayre RT (1995) Cyanogenesis in cassava (*Manihot esculenta* Crantz). *Journal of Experimental Botany*, **46**, 731–741.
- Miller RE, Tuck KL (2013) Reports on the distribution of aromatic cyanogenic glycosides in Australian tropical rainforest tree species of the Lauraceae and Sapindaceae. *Phytochemistry*, **92**, 146–152.
- Miller RE, Gleadow RM, Woodrow IE (2004) Cyanogenesis in tropical *Prunus turneri*-ana: characterisation, variation and response to low light. *Functional Plant Biology*, **31**, 491–503.
- Miller RE, Gleadow RM, Cavagnaro TR (2014) Age versus stage: does ontogeny modify the effect of phosphorus and arbuscular mycorrhizas on above- and below-ground defence in forage sorghum? *Plant Cell and Environment*, **37**, 929–942.
- Møller BL (2010) Functional diversifications of cyanogenic glucosides. *Current Opinion in Plant Biology*, **13**, 338–347.
- Montagnac JA, Davis CR, Tanumihardjo SA (2009) Nutritional value of cassava for use as a staple food and recent advances for improvement. *Comprehensive Reviews in Food Science and Food Safety*, **8**, 181–194.
- Munné-Bosch S, Alegre L (2004) Die and let live: leaf senescence contributes to plant survival under drought stress. *Functional Plant Biology*, **31**, 203–216.
- Muoki PN, Maziya-Dixon B (2010) Household utilization of Manioc (*Manihot esculenta* Crantz) in northern Mozambique. *Ecology of Food and Nutrition*, **49**, 337–356.
- Neilson EH, Goodger JQD, Woodrow IE, Møller BL (2013) Plant chemical defense: at what cost? *Trends in Plant Science*, **18**, 250–258.
- Ngudi DD, Kuo YH, Lambein F (2003) Cassava cyanogens and free amino acids in raw and cooked leaves. *Food and Chemical Toxicology*, **41**, 1193–1197.
- Nhassico D, Muquingue H, Cliff J, Cumbana A, Bradbury JH (2008) Rising African cassava production, diseases due to high cyanide intake and control measures. *Journal of the Science of Food and Agriculture*, **88**, 2043–2049.
- Niang I, Ruppel OC, Abdrabo MA, Essel A, Lennard C, Padgham J, Urquhart P (2014) Africa. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects* (eds Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Field CB, Barros VR, Dokken DJ, Mach KJ, Mastrandrea MD, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR, White LL]), pp. 1199–1265. Cambridge University Press, Cambridge, UK/New York.
- Okogbenin E, Ekanayake IJ, Porto MCM (2003) Genotypic variability in adaptation responses of selected clones of cassava to drought stress in the Sudan savanna zone of Nigeria. *Journal of Agronomy and Crop Science*, **189**, 376–389.
- Oluwole OSA (2015) Cyclical Konzo epidemics and climate variability. *Annals of Neurology*, **77**, 371–380.
- Palta JA (1984) Influence of water deficits on gas-exchange and the leaf area development of cassava cultivars. *Journal of Experimental Botany*, **35**, 1441–1449.
- Parker JD, Salminen J-P, Agrawal AA (2012) Evolutionary potential of root chemical defense: genetic correlations with shoot chemistry and plant growth. *Journal of Chemical Ecology*, **38**, 992–995.
- Pinstrup-Andersen P (2009) Food security: definition and measurement. *Food Security*, **1**, 5–7.
- Reuter DJ, Robinson JB (1997) *Plant Analysis: An Interpretation Manual*. CSIRO Publishing, Collingwood.

- Rosenthal DM, Slattery RA, Miller RE *et al.* (2012) Cassava about-FACE: greater than expected yield stimulation of cassava (*Manihot esculenta*) by future CO₂ levels. *Global Change Biology*, **18**, 2661–2675.
- Rufino MC, Thornton PK, Ng'ang'a SK, Mutie I, Jones PG, Van Wijk MT, Herrero M (2013) Transitions in agro-pastoralist systems of East Africa: impacts on food security and poverty. *Agriculture, Ecosystems and Environment*, **179**, 215–230.
- Santisopasri V, Kurotjanawong K, Chotineeranat S, Piyachomkwan K, Sriroth K, Oates CG (2001) Impact of water stress on yield and quality of cassava starch. *Industrial Crops and Products*, **13**, 115–129.
- Scott GJ, Rosegrant MW, Ringler C (2000) Global projections for root and tuber crops to the year 2020. *Food Policy*, **25**, 561–597.
- Selmar D, Kleinwächter M (2013) Stress enhances the synthesis of secondary plant products: the impact of stress-related over-reduction on the accumulation of natural products. *Plant and Cell Physiology*, **54**, 817–826.
- Setter TL, Fregene MA (2007) Recent advances in molecular breeding of cassava for improved drought stress tolerance. In: *Advances in Molecular Breeding Toward Drought and Salt Tolerant Crops* (eds Jenks MA, Hasegawa PM, Jain SM), pp. 701–711. Springer, Dordrecht, The Netherlands.
- Stochmal A, Oleszek W (1997) Changes of cyanogenic glucosides in White Clover (*Trifolium repens* L.) during the growing season. *Journal of Agricultural and Food Chemistry*, **45**, 4333–4336.
- Turyagyenda LF, Kizito EB, Ferguson M, Baguma Y, Agaba M, Harvey JJ, Osiru DS (2013) Physiological and molecular characterization of drought responses and identification of candidate tolerance genes in cassava. *AoB Plants*, **5**, Plt007.
- Van Rijssen FWJ, Morris EJ, Eloff JN (2013) Food safety: importance of composition for assessing genetically modified cassava (*Manihot esculenta* Crantz). *Journal of Agricultural and Food Chemistry*, **61**, 8333–8339.
- Vandeger R, Miller RE, Bain M, Gleadow RM, Cavagnaro TR (2013) Drought adversely affects tuber development and nutritional quality of the staple crop cassava (*Manihot esculenta* Crantz). *Functional Plant Biology*, **40**, 195–200.
- Zhao P, Liu P, Shao J *et al.* (2015) Analysis of different strategies adapted by two cassava cultivars in response to drought stress: ensuring survival or continuing growth. *Journal of Experimental Botany*, **66**, 1477–1488.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix S1. Foliar chlorophyll concentrations at final harvest, as well as methods information for the determination of mycorrhizal colonization and foliar chlorophyll concentrations, and for chlorophyll fluorescence measurements.